


FORM PTO-1390 (REV 5-93)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE ATTORNEY'S DOCKET NUMBER	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371		WARF H108	
		U.S. APPLICATION NO. (If known, 35 U.S.C. 371(c)(1)) <b>09/463510</b> Not yet assigned	
INTERNATIONAL APPLICATION NO. PCT/US98/15910	INTERNATIONAL FILING DATE 27 July 1998	PRIORITY DATE CLAIMED 30 July 1997	
TITLE OF INVENTION GERMPLASM AND MOLECULAR MARKERS FOR DISEASE RESISTANCE IN POTATO			
APPLICANT(S) FOR DO/EO/US WISCONSIN ALUMNI RESEARCH FOUNDATION			
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information			
<ol style="list-style-type: none"> <li>1. <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371.</li> <li>2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371.</li> <li>3. <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(c)(9)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).</li> <li>4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.</li> <li>5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) <ol style="list-style-type: none"> <li>a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau).</li> <li>b. <input type="checkbox"/> has been transmitted by the International Bureau.</li> <li>c. <input checked="" type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US)</li> </ol> </li> <li>6. <input checked="" type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)).</li> <li>7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) <ol style="list-style-type: none"> <li>a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau).</li> <li>b. <input type="checkbox"/> have been transmitted by the International Bureau.</li> <li>c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</li> <li>d. <input checked="" type="checkbox"/> have not been made and will not be made.</li> </ol> </li> <li>8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).</li> <li>9. <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).</li> <li>10. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).</li> </ol>			
Items 11. to 16. below concern other document(s) or information included:			
11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.			
12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.			
13. <input type="checkbox"/> A <b>FIRST</b> preliminary amendment. <input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment.			
14. <input type="checkbox"/> A substitute specification.			
15. <input type="checkbox"/> A change of power of attorney and/or address letter.			
16. <input checked="" type="checkbox"/> Other items or information: Copy of Form PCT/IB/308 (July 1996)			

09/463510 Not yet assigned		INTERNATIONAL APPLICATION NO PCT/US98/15910		ATTORNEY'S DOCKET NUMBER WARE H108	
17. <input checked="" type="checkbox"/> The following fees are submitted: <b>Basic National Fee (37 CFR 1.492(a)(1)-(5)):</b> Search Report has been prepared by the EPO or JPO.....  International preliminary examination fee paid to USPTO (37 CFR 1.482) ..... No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2))..  Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO.....  International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4).....  ENTER APPROPRIATE BASIC FEE AMOUNT = \$ 96.00				CALCULATIONS PTO USE ONLY	
Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input checked="" type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$ 130.00	
Claims	Number Filed	Number Extra	Rate		
Total Claims	15 -20 -	0	X 18	\$ 0	
Independent Claims	5 -3 -	2	X 78	\$ 156.00	
Multiple dependent claims(s) (if applicable)				+	\$
TOTAL OF ABOVE CALCULATIONS				=	\$ 382.00
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28).				\$	0
SUBTOTAL				=	\$ 382.00
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				+	\$ 0
TOTAL NATIONAL FEE				=	\$ 382.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$	0
TOTAL FEES ENCLOSED				=	\$ 382.00
				Amount to be:	
				refunded	\$
				charged	\$
a. <input checked="" type="checkbox"/> A check in the amount of \$ 382.00 to cover the above fees is enclosed. b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed. c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 04-1406. A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO: REED, Janet E. DANN, DORFMAN, HERRELL AND SKILLMAN 1601 Market Street Suite 720 Philadelphia, Pennsylvania 19103 United States of America			 SIGNATURE Janet e. Reed, Ph.D. NAME 36,252 REGISTRATION NUMBER		

09/463510  
402 Rec'd PCT/PTO 24 JAN 2000GERMPLASM AND MOLECULAR MARKERS FOR  
DISEASE RESISTANCE IN POTATO

This application claims priority to U.S. Provisional Application Serial No. 60/054,267, filed July 30, 1997, which is incorporated by reference herein in its entirety.

5 Pursuant to 35 U.S.C. §202(c), it is acknowledged that the U.S. Government has certain rights in the invention described herein, which was made in part with funds from the United States Department of Agriculture.

10

**FIELD OF THE INVENTION**

This invention relates to the field of genetic manipulation of higher plants. More specifically, the invention relates to novel germplasms, breeding stocks and molecular markers created or identified by somatic fusion of domestic and wild potato species, which are useful for development of potato varieties resistant to late blight and other fungal pathogens.

15  
**BACKGROUND OF THE INVENTION**

Several publications are referenced in this application in parentheses in order to more fully describe the state of the art to which this invention pertains. Full citations for these references are found at the end of the specification. The disclosure of each of these publications is incorporated by reference herein.

20 Plant diseases cause billions of dollars in losses to farmers in the United States and elsewhere in the world every year. Generating crop plants that are naturally resistant to disease has been a goal of plant breeders for decades. Classical breeding methods have been supplemented in recent years by molecular genetic

techniques, e.g., to identify a gene that encodes a protein with antifungal or antibacterial properties (often not a plant gene) and then express this gene at high levels in a plant.

5           Another approach is to use genes from wild species to improve disease resistance and other agronomic characteristics of cultivated crops. For the most part, however, breeders have been restricted to those genetic combinations that can be obtained by direct sexual  
10 crosses or through bridging crosses through several species.

Protoplast fusion in some cases has provided a wider range of available genes. By this technique, the somatic cells of two species are combined. Plants can  
15 then be regenerated from the combinations and examined for the expression of the desired attributes and for fertility. In this manner it is possible to incorporate useful traits from widely separated, sexually incompatible, species into breeding lines. Using other  
20 molecular techniques, it is also possible to identify genes in those wild species responsible for conferring the useful trait, such as resistance to one or more plant pathogens. The genes can then be incorporated into the genomes of a variety of different species to develop  
25 resistance to one or more plant pathogens.

Potato (*Solanum tuberosum*) is the world's fourth most valuable crop. In the United States, the value of the crop exceeds two billion dollars each year. In spite of its high value, the commercial crops are subject to  
30 many disease problems, including foliar diseases such as late blight and early blight, virus diseases, soil problems such as those caused by nematodes or *Verticillium* species, and bacterial diseases such as bacterial wilt (in the field) or *Erwinia* soft rot (in  
35 storage). These diseases are costly in terms of crop loss, expenses associated with application of chemicals and environmental impact of pesticide use. Such costs

could be minimized or avoided if resistant potato varieties were available. However, adequate resistance for late blight, *Erwinia* soft rot and many other diseases has not been incorporated into potato cultivars, partly  
5 because of the lack of a good diversity of resistance genes that breeders can use to develop resistant cultivars.

Of particular importance is the late blight disease, cause by the fungus *Phytophthora infestans*.  
10 Late blight remains one of the most devastating diseases of potatoes worldwide. Despite its importance, no major cultivars with adequate late blight resistance are grown in the United states today. Until recently, crops were protected from the disease by cultural methods (e.g. crop  
15 rotation, crop hygiene) and with fungicides. The absence of compatible mating types within the U.S. heretofore has prevented sexual recombination. However, a second mating type has now become widespread in the U.S. and many lines of the fungus have become resistant to one of the key,  
20 very effective, systemic fungicides (Metalaxyl) registered for potato late blight control.

The late blight fungus is also a devastating pathogen on crops other than potato. It infects tomatoes, eggplants and other solanaceous species. Other  
25 *Phytophthora* species are pathogenic to a wide array of agronomically important plants, including grapes, avocados and several varieties of fruit and nut trees. Accordingly, a source of resistance to *Phytophthora* species that could be introduced into these species by  
30 molecular genetic techniques would also be of great value.

Possible sources of resistance to many potato pathogens exist in related *Solanum* species. Several *Solanum* species have been crossed with the cultivated  
35 potato in an effort to introgress disease-resistance genes.

When sexual crossing techniques have failed,

disease resistance transfers have been attempted with somatic cell fusions. Potato protoplasts have been fused with a number of sexually incompatible wild *Solanum* species (e.g., *S. brevidens*, *S. bulbocastanum*, *S. commersonii*, *S. polyadenium*, *S. etuberosum*), and many fertile somatic hybrids have been regenerated (Austin et al., 1985, 1993; Ehlenfeldt & Helgeson, 1987; Kim-Lee et al., 1993; Novy & Helgeson, 1994b). Somatic hybrids have been screened for useful disease resistance (Helgeson et al., 1986; Austin et al., 1988; Novy & Helgeson, 1994b) and these resistances are heritable (Helgeson et al., 1993).

*Solanum bulbocastanum* is a particularly desirable wild species from which to seek useful disease resistance genetic traits, inasmuch as it exhibits resistance to several potato pathogens, including nematodes, early blight, late blight and *Verticillium*. Disease- or pest-resistant somatic hybrids of *S. bulbocastanum* and cultivated potato have been produced by the present inventors and by others. In one instance, analysis of BC1 and BC2 progeny of a nematode-resistant *S. bulbocastanum* - *S. tuberosum* somatic hybrid revealed that the nematode resistance locus is likely to reside on chromosome 11 of *S. bulbocastanum* (Brown et al., 1996). However, the chromosomal location of late blight resistance in *S. bulbocastanum* heretofore has not been identified.

Clearly, there is an ongoing need to identify such resistances in wild potato species, such as *S. bulbocastanum*, to continue improving the disease resistant characteristics of cultivated potatoes. A need also exists to identify the genes in wild potato species that are responsible for disease resistance. Once isolated, these genes can then be introduced by molecular genetic techniques into species other than potato to confer resistance to one or more plant pathogens.

**SUMMARY OF THE INVENTION**

This invention provides novel germplasms, breeding stocks, molecular markers and methods for introducing late blight resistance into cultivated potato plants. The invention further provides genomic DNA segments from *S. bulbocastanum* useful for introducing resistance to the late blight fungus, *Phytophthora infestans*, into species other than potato.

According to one aspect of the invention, a potato germplasm is provided that confers resistance to the late blight fungus, *Phytophthora infestans*, as well as other fungal pathogens, including early blight, *Erwinia* soft rot and *Verticillium*. The most fundamental form of this germplasm is a tissue culture produced by somatic hybridization of *S. tuberosum* with *S. bulbocastanum*. Fertile plants regenerated from these hybrids are also provided, along with progeny resulting from crosses with agronomically preferable cultivated potato species.

According to another aspect of the invention, a late blight-resistant potato plant is provided, comprising a segment of a genome from *Solanum bulbocastanum* which contains a gene that confers resistance to late blight. In a preferred embodiment, the genomic segment of *S. bulbocastanum* is from chromosome 8, and co-segregates with one or more of the following markers: (1) a RAPD marker referred to herein as GO2<sub>586</sub>; (2) a RAPD marker referred to herein as PO9<sub>587</sub>; and (3) RFLP marker CT88, RFLP marker, RFLP marker CT148, RFLP marker CT252 and RFLP marker CT68. The potato plant may also be resistant to at least one additional disease, such as potato early blight, *Erwinia* soft rot, and *Verticillium* wilt.

In a preferred embodiment, the aforementioned late blight resistance gene is incorporated into the potato plant by somatic hybridization between a cell of a parent of the plant and a cell of *Solanum bulbocastanum*.

In another embodiment, the late blight resistance gene is incorporated into the plant by genetic transformation of a cell of the plant with a plant transformation vector containing the gene.

5           According to another aspect of the invention, an isolated nucleic acid molecule is provided, which is complementary to part or all of a 0.6 kb segment of a *Solanum bulbocastanum* genome, which co-segregates with a gene that confers resistance to late blight. In a  
10 preferred embodiment, the segment comprises part or all of the RAPD marker GO2<sub>586</sub> or PO9<sub>587</sub>, having the sequence of SEQ ID NO:1 or SEQ ID NO:2, respectively.

          According to another aspect of the invention, a method of monitoring late blight resistance in a breeding  
15 cross of progeny of a fertile somatic hybrid of *Solanum tuberosum* and *Solanum bulbocastanum* is provided. The method comprises: (a) performing the cross; (b) isolating genomic DNA from individual progeny of the cross; and (c) detecting in the genomic DNA the presence  
20 or absence of a genetic marker that is pre-determined to co-segregate with the late blight resistance, the presence or absence of the marker being indicative of the presence or absence of the late blight resistance in the individual progeny of the breeding cross. In a preferred  
25 embodiment, the marker is selected from the group of RAPD and RFLP markers listed above.

          According to yet another aspect of the invention, a method of identifying a *Solanum*  
*bulbocastanum* gene that confers resistance to late blight  
30 is provided. The method comprises: (a) cloning a DNA segment that co-segregates with the late blight resistance phenotype in progeny of somatic hybrids of *Solanum bulbocastanum* and *Solanum tuberosum*; (b) providing a genomic library of the *Solanum bulbocastanum*  
35 genome; (c) isolating clones of the genomic library that contain segments which hybridize with the co-segregating DNA segment; and (d) identifying at least one gene



disposed within the isolated genomic clones that confers the late blight resistance. In a preferred embodiment, the cloned DNA segment that co-segregates with late blight resistance comprises part or all of one of the

5 RAPD or RFLP markers listed above.

According to further aspects of the invention, a late blight resistance gene from *Solanum bulbocastanum*, produced by the aforementioned method, is provided. Also provided is a transgenic plant comprising the resistance

10 gene.

The advantages of the present invention with be better understood in view of the detailed description and examples set forth below.

#### 15 BRIEF DESCRIPTION OF THE DRAWING

Figure 1. RFLP analysis of somatic hybrids between *S. bulbocastanum* and potato. The probe used in the analysis was TG310, a tomato genomic probe specific for Chromosome 1 of tomato and potato.

20 Figure 2. MapMaker analysis of chromosome 8 of *Solanum bulbocastanum* with RAPD (randomly amplified polymorphic DNA) and RFLP (restriction fragment length polymorphism) markers and resistance to late blight. Percentages in parentheses (left column) indicate recombination frequencies calculated by dividing the

25 deviation from complete co-inheritance by the population size. Numbers to the right of the percentage column indicate distances between markers in centimorgans. Numbers in parentheses to the right of the chromosome 8

30 diagram represent the arbitrary code number of the individual in the population. The far right column lists RAPD markers, which are named by the decameric primer and the size of the amplified transcript (e.g. "GO2-586" or "PO9-587") and RFLP markers. The resistance locus is

35 indicated by "R". Map scale is 10.0 cM per 1.21 cm.

Figure 3. Tomato chromosome 8 map, after Tanksley et al., 1992, showing approximate corresponding

location of *S. bulbocastanum* late blight resistance gene(s) on tomato chromosome 8.

#### DETAILED DESCRIPTION OF THE INVENTION

5           In accordance with the present invention, genetic markers have been identified that are linked to late blight resistance derived from somatic hybrids of *Solanum bulbocastanum* and cultivated potato (*Solanum tuberosum*). Backcross (BC<sub>2</sub>) populations were analyzed for  
10 resistance to late blight and for randomly amplified polymorphic DNA (RAPD) and RFLP markers. Three test populations were derived from two different somatic hybrids between potato and *S. bulbocastanum*. Each somatic hybrid was crossed with the cultivar Katahdin to  
15 generate BC<sub>1</sub> parents, all of which were resistant to late blight. BC<sub>1</sub> progeny were crossed with three different potato breeding lines (Norland, Atlantic and A 89804-7), all of which are sensitive to the late blight fungus. Each BC<sub>2</sub> population contained more than 50 individuals and  
20 segregated for resistance to late blight. In each population, late blight resistance was correlated (>95%) with the presence of a RAPD marker ("GO2<sub>586</sub>") keyed to chromosome 8 of *S. bulbocastanum*. The identification of this novel molecular marker is described in greater  
25 detail in Example 2.

          Further genetic analysis has led to the identification of another RAPD marker and several RFLP markers that are also associated with the resistance gene or genes of *S. bulbocastanum* chromosome 8. These  
30 include, for example, RFLP marker CT88 and RAPD marker PO9<sub>587</sub>, which appear to flank the resistance locus (see Figure 2). Other closely linked RFLPs include CT148, CT252 and CT68 (Figure 2). Together, these markers define a specific location on *S. bulbocastanum* chromosome  
35 8 that carries the gene(s) which confer resistance to late blight and certain other potato diseases, as discussed in greater detail herein. The approximate

location of the resistance genes on Chromosome 8 (as mapped by Tanksley et al., 1992) is shown in Figure 3.

This invention provides a new and useful germplasm and breeding stock for breeding potato cultivars with resistance to late blight. The germplasm comprises fertile hybrids produced by somatic fusion of *S. bulbocastanum* with *S. tuberosum*, and progeny thereof, which contain that portion of the *S. bulbocastanum* genome carrying the late blight resistance gene or genes. The presence of this genomic fragment is conveniently monitored by the presence of closely linked RAPD markers, GO2<sub>586</sub> or PO9<sub>587</sub>, or closely-linked RFLP markers, such as CT88, as described in greater detail below. Particularly preferred breeding stock is obtained by repeated backcrosses of the somatic hybrid with potato cultivars having desirable agronomic qualities, with the presence of the late blight resistance-conferring genomic segment of *S. bulbocastanum* being monitored by detection of one or more of the relevant RAPD or RFLP markers.

Creation of *S. tuberosum* - *S. bulbocastanum* somatic hybrids, selection of fertile, resistant plants, and production of subsequent backcross generations comprising the disease resistance gene or genes are all accomplished by methods well known to plant breeders and molecular biologists. Preferred methods are described in greater detail in Example 1.

This invention also provides novel molecular markers to facilitate selection of breeding progeny that contain the resistance-conferring genomic segment from *S. bulbocastanum*, without having to perform field or greenhouse trials for disease resistance. One closely linked marker, GO2<sub>586</sub>, was created through the use of RAPD markers, using commercially available oligonucleotide 10-mers as primers for PCR amplification. RAPD markers are generated by incubating genomic DNA with a population of 10-mers under conditions that allow the oligonucleotides to bind to any complementary sequences in the genomic

DNA. The length of DNA between two bound sets of primers is amplified by PCR, thereby generating a DNA segment which is a copy of the segment between the primers in the genomic DNA, i.e. a RAPD fragment. The nucleotide  
5 sequence of the GO2<sub>586</sub> RAPD marker is set forth herein as SEQ ID NO:1.

RAPD fragments thus created are species specific markers which can be keyed to particular chromosomes by comparative RFLP analysis and can be  
10 followed as dominant markers through various crosses. In accordance with the present invention, RAPD marker GO2<sub>586</sub> (which is a 586 bp fragment primed by the "GO2" decameric oligonucleotide purchased from Operon Technologies, Inc.) was keyed to *S. bulbocastanum* chromosome 8. Analysis of  
15 BC<sub>2</sub> progeny of potato-*S. bulbocastanum* somatic hybrids demonstrated that late blight resistance correlates with greater than 95% frequency with the presence of the GO2<sub>586</sub> RAPD fragment. This tight segregation of late blight resistance with the marker indicates that the resistance  
20 gene or genes reside at or near the location of the marker. Accordingly, late blight resistance in further crosses can be monitored by detecting the presence or absence of the marker. In this manner, progeny having a high probability of carrying the resistance gene(s) can  
25 be selected without lengthy greenhouse or field trials for disease resistance, thus making the breeding process faster and more efficient.

This invention also provides a second RAPD marker, PO9<sub>587</sub>, which is also closely linked with the  
30 resistance gene(s) in *S. bulbocastanum*, and which was identified by the same protocol as set forth above for GO2<sub>586</sub>. The nucleotide sequence of RAPD marker PO9<sub>587</sub> is set forth herein as SEQ ID NO:2.

This invention further provides RFLP molecular  
35 markers, useful in facilitating selection of breeding progeny that contain the resistance-conferring segment of *S. bulbocastanum*. These markers can also assist in

defining the location of the resistance genes on the chromosome, and obtaining isolated genomic segments containing the gene(s). The nucleotide sequence of RFLP CT88 from three different sources (published by Tanksley et al. (<http://probe.nalusda.gov:8300/cgi-bin/browse/solgenes>), from R4 potato, and from *S. bulbocastanum*) are set forth herein as SEQ ID NOS: 3, 4 and 5 respectively).

Once created, useful RAPD or RFLP fragments can be maintained in any suitable cloning vector. For instance, the GO2<sub>586</sub> marker generated in accordance with the present invention is maintained in a plasmid vector provided with a commercially available PCR kit (Invitrogen, Inc.). It is noted that obtaining RAPD should be replicable by anyone of skill in the art, using the commercially available decamers and the methods described in Example 2 and references cited therein. Alternatively, RAPD fragments described herein can be duplicated by nucleotide synthesis using standard methodologies.

The RAPD fragments, or portions thereof, or any of the linked RFLPs discussed above or shown in Figure 2, are used to monitor the presence or absence of the late blight resistance gene(s) by labeling them as probes to hybridize with complementary sequences in the *S. bulbocastanum* genome. The cloned fragments may be labeled according to any standard means, many of which are set forth in "Current Protocols in Molecular Biology", eds. Frederick M. Ausubel et al., John Wiley & Sons, 1997. The complementary genomic DNA is detected by one of several standard methods including, but not limited to, (1) *in situ* hybridization; (2) Southern hybridization (3) "dot blot" hybridization; and (4) assorted amplification reactions such as polymerase chain reactions (PCR). Detection of the complementary genomic DNA is indicative of the presence of the late blight resistance-conferring gene(s), due to the close linkage

of the R gene(s) and the RAPD marker or RFLPs, as discussed above.

The RAPD fragments, or portions thereof, or any of the aforementioned RFLP fragments, may also be used to  
5 identify and isolate the closely-linked late blight resistance gene of *S. bulbocastanum*, using methods well known to molecular biologists. In a preferred embodiment, a genomic library of *S. bulbocastanum* is constructed a suitable cloning vehicle, e.g. cosmid,  
10 yeast artificial chromosome (YAC) or bacterial artificial chromosome (BAC). The library is then screened by hybridization with the cloned RAPD fragment and/or one or more of the RFLP markers, and hybridizing clones are isolated. The hybridizing clones are then further  
15 analyzed, e.g. by mapping, sequencing and transcript analysis, to identify candidate open reading frames that may encode the resistance factor(s). Once candidate open reading frames are identified, they in turn may be further analyzed (e.g. by construction and *in vitro*  
20 expression of a cDNA molecule) in order to characterize the resistance gene and its encoded protein.

The disease resistance-conferring clone thus identified may be used to introduce late blight resistance into cultivated potato by molecular genetic  
25 techniques. For example, a binary bacterial artificial chromosome (BIBAC) vector may be used to mobilize a BAC genomic insert into potato (the vector BIBAC2 (Hamilton et al, 1996) has been used to mobilize large (>150 kb) DNA inserts into tobacco). A BIBAC vector containing the  
30 relevant genomic insert from *S. bulbocastanum* can be transferred into an *Agrobacterium tumefaciens* and used to transform selected potato cultivars. Alternatively, potato cells can be transformed via biolistic delivery of the BIBAC clone. Putative transgenic clones can then be  
35 evaluated by standard methods to determine if the transformation has been successful.

The disease resistance-conferring clone also

may be used to introduce resistance to *Phytophthora infestans* in species other than potato, which are sensitive to infection by that organism. These species include, but are not limited to, tomato, eggplant and other *Solanum* species. Furthermore, it is possible that the late blight resistance gene(s) of *S. bulbocastanum* may confer resistance to diseases other than late blight, and so may be of even broader utility for introducing disease resistance in potato and other plant species. The gene may be particularly useful for introducing resistance to other *Phytophthora* species pathogenic to grapes, avocados, fruit trees and nut trees. Moreover, once the gene's function has been determined, this may lead to the discernment of new mechanisms of resistance in other species.

The following examples are provided to describe the invention in greater detail. They are intended to exemplify, not to limit, the scope of this invention.

#### EXAMPLE 1

##### Resistance to Late Blight in Somatic Hybrids of *Solanum bulbocastanum* and Potato, and Progeny Thereof

A Mexican wild species, *S. bulbocastanum*, is highly resistant to late blight. However, *S. bulbocastanum* is a 1EBN species and thus extremely difficult to cross directly with potato.

Somatic hybridization can provide a means of bypassing sexual incompatibility between *Solanum* species, leading to fertile plants that can be used directly in breeding programs. The experimental results set forth in this Example demonstrate that the resistance in *S. bulbocastanum* can be captured and passed on to potato breeding lines by the use of somatic hybridization.

#### MATERIALS AND METHODS

Potato and related species used for somatic hybridization were obtained from Dr. John Bamberg and his

colleagues at the Inter-Regional Potato Introduction Station (NRSP-6), 4312 Highway 42, Sturgeon Bay, WI. These include *S. bulbocastanum*, PI 243510 and *S. tuberosum* PI 23900 (potato). Elite copies of potato  
5 cultivars (Katahdin, Atlantic) were obtained from the Wisconsin potato certification program. All cultivars and wild species, and test materials were routinely maintained clonally *in vitro* as described by Haberlach et al (1985). The individual clones were multiplied *in*  
10 *vitro* for analyses.

Protoplasts were isolated from leaves of *in vitro* shoots as described by Haberlach et al. (1985). Somatic hybridizations with the protoplasts were performed using a polyethylene glycol (PEG) protocol.  
15 For the most part, the procedure of Austin et al. (1985) was followed. However, after PEG additions, dilutions and pelleting of the cells after the fusion attempts, the cells were suspended in 0.3 M sucrose rather than 0.6 M mannitol. The cell suspension was gently shaken (40 RPM)  
20 for 45 minutes and then centrifuged (HNII centrifuge, IEC) at 1300 rpm for 10 minutes in a Babcock bottle. This modification resulted in viable protoplasts and fused cells being concentrated at the surface of the sucrose solution in the bottle, thus separating the  
25 viable cells from the pelleted debris.

The resulting fused cells were regenerated into whole plants in a manner similar to that reported by Haberlach et al. (1985). Initially, the cells were plated onto culture medium (CUL, Haberlach et al. 1985)  
30 and, after macroscopic calli had appeared, the calli were transferred to differentiation medium (DIF, Haberlach et al. 1985). After 2-3 weeks, the calli were transferred to the differentiation medium developed by Lam (1977). After buds had formed, the calli were  
35 transferred to proliferation medium (PM, Haberlach et al. 1985). Shoots that formed were then excised and rooted on standard propagation medium (PROP, Haberlach et al.



1985) and maintained in test tubes *in vitro*. Clonal copies of the reference copy were made for experiments.

DNA extractions and restriction fragment length polymorphism (RFLP) analyses were carried out as described by Williams et al. (1990). Chromosome specific tomato genomic (TG) and cDNA (CD) probes were obtained from Dr. Steve Tanksley, Cornell Univ. Four putative somatic hybrids were analyzed by this method. To complete the analyses for hybridity, randomly amplified polymorphic DNA (RAPD) analyses were carried out as described in greater detail in Example 2. A total of 109 primers (from 380 primers tested) were selected that give clearly scorable polymorphisms between potato and *S. bulbocastanum*. Several of these were used with each of the putative somatic hybrids.

Three of the hybrids have been used extensively in further experiments. These were designated J101, J103, and J138. In crosses, the potato parent was designated with K (for Katahdin) or A (for Atlantic). Thus, for example, J101K1 was the first seed germinated from a berry obtained from the cross of J101 and Katahdin. Similarly, J101K6 and J101K27 were the seedlings obtained from seed 6 and seed 27 respectively from that cross. The BC2 progeny were named by adding the seed number and cultivar lines in that cross. Thus the cross designated as J101K6A22 is the 22nd seedling from the cross of line J101K6 with Atlantic. This shortcut avoided use of the longer and less informational term ((*S. bulbocastanum* + *S. tuberosum*) x Katahdin) x Atlantic that could be applied to this individual.

Comparisons were made of susceptible and resistant plants in the field. For these studies, the percentage of leaves showing late blight lesions was recorded at various times during the growing season. For detailed greenhouse studies, a modified Horsfall-Barret rating scheme was used to estimate percent leaf infection by *P. infestans*. The ratings and the ranges of %

infections associated with these scores are as follows:  
9, no visible infection; 8, <10%; 7, 11-25%; 6, 26-40%;  
5, 41-60%; 4, 61-70%; 3, 71-80%; 2, 81-90% 1, >90%;  
0, 100% infection.

5

## RESULTS

In total, 80 shoots were obtained from 23  
calli. Twenty four of these plants (from 5 calli) showed  
obvious morphological differences from either of the  
parent species. The other 56 plants appeared very  
similar to potato. Initially, chromosome-specific  
restriction fragment length polymorphism (RFLP) markers  
were used to confirm that four of the plants derived  
from fusion of *S. bulbocastanum* and potato cells were  
indeed somatic hybrids. Prominent potato bands were  
retained by the hybrids in addition to the diagnostic *S.*  
*bulbocastanum* bands (Figure 1). The rest of the  
potentially hybrid plants were evaluated with RAPD probes  
(see Example 2). A total of 13 more somatic hybrids  
were confirmed by these techniques. The confirmed  
hybrids were derived from four different callus pieces,  
and thus probably from four different fusion events.

The appearance of leaves and stems of the  
parent plants and representative somatic hybrids was  
examined. As has been the case with many of our other  
hybrids, traits of both of the parent species could be  
seen in the hybrids. In this case, the purple coloration  
of the *S. bulbocastanum* stems was expressed in the  
hybrids. However, leaves of the hybrids were compound  
rather than single as was the case of the wild species.

Crosses of four of the somatic hybrids were  
undertaken with the potato cultivars "Katahdin" and  
"Atlantic" to test for fertility of the hybrids. Each of  
the tested hybrids yielded viable seeds and sexual  
progeny. Further crosses of selected individuals from  
these progeny lines were also successful. Thus, parental  
lines and two successive backcross populations were

available for evaluation of disease resistance.

Preliminary laboratory tests for resistance to *P. infestans* were made with detached leaves or leaf discs indicated that the somatic hybrids and some of the  
5 progeny retained at least some of the resistance to late blight that was shown by the *S. bulbocastanum* parent.

These preliminary laboratory results were confirmed in field tests during a first growing season. Somatic hybrids as well as progeny from somatic hybrids  
10 between *S. bulbocastanum* and potato, showed remarkable resistance and were clearly noticeable in test plots as "green islands " in a brown background of dead potato lines. Although the cultivars Atlantic, Russet Burbank, and Snowden were killed, eleven different experimental  
15 lines showed less than 10% infection. In each case, the live test plants were surrounded with the susceptible cultivar, Russet Burbank, which had been killed by the fungus by August 9 of that growing season. In contrast, % foliage infections on August 15 were 5.0% and 7.8% for  
20 J101K27 and J101K6A22, respectively.

Fourteen BC<sub>1</sub> and BC<sub>2</sub> lines were tested in Toluca, Mexico in a second growing season the following summer. Good resistance in the Toluca field test was also obtained with all lines that were resistant in  
25 Wisconsin in the previous summer.

Additional field experiments were carried out at Hancock, Wisconsin, in a third growing season. Again, a severe natural late blight epidemic was obtained in test plots and yields of common cultivars were severely  
30 depressed by the late blight epidemic. For example, in plots where an effective fungicide spray regime was maintained, Russet Burbank yields were as high as 1.7 kg/plant. In the yield trial where no fungicide was used, this was cut almost in half to 0.86 kg/plant. One  
35 of the *S. bulbocastanum* -derived lines , J103K7, topped all 90 test lines with a yield of 1.36 kg/plant and J138A12 ranked fourth at Hancock in the third growing

season, with a yield of 1.32 kg/plant.

To test the resistance of potentially segregating BC<sub>1</sub> and BC<sub>2</sub> populations, a facility was constructed in the new research greenhouses at the University of Wisconsin Biotron. There, close control of humidity and temperature made uniform epidemics possible. Segregation of resistance and susceptibility was obtained for each of six BC<sub>1</sub> from four different somatic hybrids.

Representative tests from one of these lines are shown in Table 1. Three of these lines were further crossed to Atlantic or Norland. Representative results on these BC<sub>2</sub> lines are included in Table 2. Again, clear segregation of resistance and recovery of both parental extremes of susceptibility and resistance were obtained with these lines.

**Table 1. Representative data from Biotron tests of BC<sub>1</sub> lines for late blight resistance**

20	Plant line	Average Blight score		
		5day	8 day	12 day
25	J101K09	9.0 ± 0.0	9.0 ± 0.0	9.0 ± 0.0
	J101K27	9.0 ± 0.0	9.0 ± 0.0	9.0 ± 0.0
	J101K06	8.8 ± 0.5	9.0 ± 0.0	8.8 ± 0.4
	J101K10	9.0 ± 0.4	8.8 ± 0.4	8.6 ± 0.9
	J101K30	9.0 ± 0.0	9.0 ± 0.0	8.4 ± 0.9
30	J101K16	8.8 ± 1.0	8.8 ± 0.5	7.8 ± 1.0
	J101K25	8.6 ± 1.1	8.4 ± 0.9	8.2 ± 1.1
	J101K02	8.6 ± 0.4	8.0 ± 1.2	7.8 ± 0.4
	J101K20	8.6 ± 0.7	7.4 ± 0.9	7.0 ± 0.7
	J101K33	7.4 ± 1.8	5.4 ± 0.9	6.2 ± 1.8
35	J101K19	6.8 ± 2.4	5.6 ± 2.7	5.2 ± 2.4
	J101K11	6.0 ± 1.3	4.6 ± 1.1	4.2 ± 1.3
	J101K12	7.2 ± 2.5	5.2 ± 2.4	3.8 ± 2.5
	J101K18	6.8 ± 1.8	4.6 ± 1.3	3.6 ± 1.8
	J101K21	5.2 ± 1.3	2.2 ± 1.3	0.6 ± 1.3
40	S. bulbocastanum	9.0 ± 0.0	9.0 ± 0.0	9.0 ± 0.0
	Somatic hybrid J101	8.6 ± 0.5	7.8 ± 0.8	8.2 ± 0.0
	S. tuberosum			
	PI 203900	7.0 ± 0.0	5.8 ± 1.3	5.6 ± 1.9
	S. tuberosum			
45	cv "Kathadin"	4.8 ± 0.4	2.0 ± 0.7	1.0 ± 1.2

Table 2. Examples of segregation for late blight resistance in BC<sub>2</sub> progeny of a cross between BC<sub>1</sub> line J101K6 and *S. tuberosum* cv. Atlantic.

5	Plant line	Average blight score		
		7 day	10day	15 day
10	J101K6A21	9.0 ± 0.0	9.0 ± 0.0	9.0 ± 0.0
	J101K6A4	8.8 ± 0.4	8.8 ± 0.4	9.0 ± 0.0
	J101K6A22	9.0 ± 0.0	9.0 ± 0.0	8.8 ± 0.4
	J101K6A2	9.0 ± 0.0	9.0 ± 0.0	8.8 ± 0.4
	J101K6A3	5.2 ± 3.1	5.4 ± 3.6	2.0 ± 3.9
15	J101K6A10	3.4 ± 2.1	3.0 ± 2.0	1.4 ± 1.9
	J101K6A50	5.6 ± 3.4	3.4 ± 3.6	0.0 ± 0.4
	J101K6A24	2.6 ± 0.9	1.4 ± 0.5	0.0 ± 0.0
	<i>S. bulbocastanum</i>			
	PI 243510	9.0 ± 0.0	8.8 ± 0.4	9.0 ± 0.0
20	<i>S. tuberosum</i>			
	PI 203900	4.0 ± 1.0	4.0 ± 1.0	0.6 ± 0.9
	J101*	7.8 ± 1.1	8.6 ± 0.5	8.4 ± 0.9
	Kathadin*	4.4 ± 0.5	4.2 ± 0.8	3.4 ± 1.8
	J101K6**	9.0 ± 0.0	9.0 ± 0.0	9.0 ± 0.0
25	Atlantic**	3.6 ± 0.5	3.0 ± 0.0	1.8 ± 1.8
	* Lines crossed to generate BC <sub>1</sub> lines			
	** Lines crossed to generate BC <sub>2</sub> lines.			

These results indicate that somatic hybrids between *S. bulbocastanum* and potato are sources of highly effective resistance against late blight. Furthermore, this trait can be passed on to potato breeding lines by conventional sexual crossings. The resistance carries over at least two generations of sexual progeny and, with the third season results just obtained, has been stable for 4 different years in several different locations. As no North American cultivar currently has adequate resistance against this disease, these lines will be very useful for introducing resistance into commercial lines.

The resistance to *P. infestans* from *S. bulbocastanum* appears to be more general than the race-specific resistance derived from *S. demissum* (Black & Gallegly, 1957). Nearly every race of the fungus is known to be found in Toluca, Mexico and was actually isolated from the fields where the progeny of the somatic hybrids showed good resistance. Observations of the

foliage in Toluca in two different growing seasons indicated that some lesions actually formed and that limited sporulation also occurred. Although the disease resistance is highly effective, it is unclear, as yet, the numbers of genes involved. For each of the somatic hybrids tested, the disease resistances of BC<sub>1</sub> lines appear to segregate. Thus, it appears that the late blight resistance gene(s) is (are) heterozygous in the clone of *S. bulbocastanum* utilized in somatic hybridization.

#### EXAMPLE 2

##### Identification of RFLPs and a RAPD Marker Linked to Late Blight Resistance Derived from *Solanum bulbocastanum*-Potato Somatic Hybrids

Example 1 described the production of potato-*S. bulbocastanum* somatic hybrids and their progeny, which are resistant to potato late blight. This example describes the identification of DNA markers that co-segregate with disease resistance in these progeny.

Detailed genetic mapping of traits in potato has become possible due to the completion by Bonierbale et al. (1988) of a restriction fragment length polymorphism (RFLP) map in potato. By using tomato clones that had been mapped previously in tomato, a potato map was constructed via a diploid, interspecific cross. The tomato map and the potato map exhibit a nearly collinear gene order (Tanksley et al., 1992).

An alternative method for generating markers is the use of randomly amplified polymorphic DNAs (RAPDs). This technique uses DNA polymerase, synthetic oligonucleotides (10-mers) and genomic DNA mixed together in a PCR thermal cycler to generate bands that are copies of sequences of the genomic DNA present in the mix (J.G.K. Williams et al., 1990). The procedure usually generates between about five and eight specific bands that can be followed, as dominant markers, through various crosses. Moreover, if specific bands can be

linked to a characteristic (such as disease resistance) or to a particular chromosome, or both, the bands can be excised and amplified, then used as a standard RFLP. The method works well to evaluate the extension of DNA introgression from a wild species into a cultivated species. However, RAPDs are species specific, so it is necessary to develop a set for each different species, unlike with RFLPs where inter-species synteny applies.

## 10 MATERIALS AND METHODS

Somatic hybrids *S. bulbocastanum* (PI 243510) and *S. tuberosum* (PI 203900) were generated by the method of Austin et al. (1985 and 1993), as describe in detail in Example 1. Plants were derived from fertile hybrids, and three lines of these were designated J101, J103 and J138, respectively. The somatic hybrid plants were crossed as the female parent with *S. tuberosum* cv Katahdin (KAT) to give BC<sub>1</sub> progeny. Three BC<sub>1</sub> progeny were crossed as seed parents with three different potato breeding lines (Norland, Atlantic and A 89804-7) to generate BC<sub>2</sub> populations.

For RFLP and RAPD analysis, DNA was isolated exclusively from plants maintained in axenic *in vitro* cultures. DNA manipulations, PCR amplification protocols for RAPD markers (Williams et al, 1990) and assignment of RAPD markers to *S. bulbocastanum* chromosomes by reference to RFLP markers was performed as described by McGrath et al. (1994), with the exception that a modified thermocycle profile was used, as described by McGrath et al. (1996). RAPD markers were named by the deca-nucleotide primer (obtained from Operon Technologies) and the size of the amplified fragment in subscript (e.g., a 586 bp fragment amplified by primer GO2 is represented as GO2<sub>586</sub>).

Segregation of RAPD markers was also analyzed with maximum likelihood algorithms contained in the MapMaker computer package (Lander et al., 1987).

MapMaker version 2.0 for the MacIntosh was used. The data were coded and analyzed as a "Haploid" population under the "Data Type" option. The use of MapMaker for the analysis of interspecific somatic hybrids is non-standard and does not provide three-point linkage data. However, recombination frequencies do have meaning in this context. Markers that show identical segregation have a recombination frequency of 0.0%. Markers that deviate from complete co-inheritance by a single change (e.g. a marker is present in one additional individual) show a recombination frequency proportional to the population size; in this instance 1/101 individuals or 1.0%. Thus, multiples of this value indicate the number of differences observed between any pair of markers.

## RESULTS

Using RAPD marker analysis, twelve synteny groups were established for *S. bulbocastanum*, corresponding to the base number of chromosomes in that species. The groups were associated with chromosomes by comparative RFLP analysis. A MapMaker analysis of RFLPs and RAPD fragments of synteny group A (chromosome 8) of *S. bulbocastanum* is shown in Figure 2. This synteny group contains the RAPD markers GO2<sub>586</sub> and PO9<sub>587</sub> as well as the RFLP CT88. It appears that the resistance area (R) is flanked by CT88 and GO2<sub>586</sub> on one side and PO9<sub>587</sub> on the other (Figure 2).

The nucleotide sequence of GO2<sub>586</sub> is set forth herein as SEQ ID NO: 1. The nucleotide sequence of PO9<sub>587</sub> is set forth herein as SEQ ID NO:2. Three nucleotide sequences of CT88 are set forth herein. SEQ ID NO:3 is the sequence published by Tanksley et al. (<http://probe.nalusda.gov.8300/cgi-bin/browse/solgenes>); SEQ ID NO:4 is from R4 potato, and SEQ ID NO:5 is from *S. bulbocastanum* (PT29). Slight differences were noted among the three sequences. The R4 potato marker is 589 bp in length, while the *S. bulbocastanum* RFLP is 592 bp



and the Tanksley et al. sequence is 596 bp. In addition, the *S. bulbocastanum* CT88 homolog possesses two TaqI sites, whereas the other two have only one.

The fusion of *S. tuberosum* with *S.*

5 *bulbocastanum* yielded 17 confirmed somatic hybrids. The somatic hybrids are quite resistant to early blight and to late blight. The progeny of some crosses were found to segregate for high resistance to both early blight and late blight. Some of the progeny are highly resistant  
10 even to fungal lines that are highly virulent, complex races. Several highly resistant clones were selected and further crossed with potato. Three sets of BC<sub>2</sub> mapping populations have been generated by crosses with potato cultivars. Field studies in Wisconsin first indicated  
15 resistance to *Verticillium*, early blight and late blight in these materials. Subsequent studies on late blight resistance were carried out in North Dakota, Prince Edward Island, Washington state, New York, Maine, West Virginia and Toluca, Mexico. The Mexico and Wisconsin  
20 studies were repeated for a second and third season and late blight resistance in these materials appears to be durable.

Segregation of late blight resistance in selected BC<sub>1</sub> and BC<sub>2</sub> populations were shown in Example 1,  
25 Tables 1 and 2. Table 3 below shows results of segregation analysis of BC<sub>2</sub> progeny with respect to late blight resistance in relation to the presence or absence of RAPD marker GO2<sub>586</sub> for Chromosome 8 of *S. bulbocastanum*. The data shown in Table 3 were generated  
30 by PCR amplification, followed by agarose gel electrophoresis and ethidium bromide staining to observe the presence or absence of an amplified 586 bp band.

Table 3. Segregation of BC<sub>2</sub> progeny in relation to the presence or absence of RAPD marker GO2<sub>586</sub>.

5	<u>Clone</u>	<u>Late blight rating</u>			<u>Marker</u>
		<u>7 day</u>	<u>10 day</u>	<u>15 day</u>	
	PI 245310 (BLB)	9.0	8.8	9.0	+
	PI 203900 (TBR)	4.0	4.0	0.6	-
10	J101	7.8	8.6	8.4	+
	Kathadin	4.4	4.2	3.4	-
	J101K6	9.0	9.0	9.0	+
	Atlantic	3.6	3.0	1.8	-
	J101K6A22	9.0	9.0	8.8	+
15	J101K6A50	5.6	3.4	0.0	-
	J101K6A32	8.6	8.4	9.0	+
	J101K6A38	2.8	3.4	1.4	-
	J101K6A21	9.0	9.0	9.0	+
	J101K6A07	6.2	7.0	5.3	-
20	J101K6A12	9.0	9.0	9.0	+
	J101K6A03	5.2	5.4	2.0	-
	J101K6A19	9.0	9.0	8.8	+
	J101K6A18	7.0	7.7	5.3	-
	J101K6A02	9.0	9.0	8.8	+
25	J101K6A54	5.0	3.2	1.0	-

As can be seen from Table 3 (as well as other results not shown), late blight resistance in the BC<sub>2</sub> clones is highly correlated (>95%) with the presence of RAPD marker GO2<sub>586</sub>, which has been keyed to chromosome 8 of *S. bulbocastanum*. This high correlation of the resistance phenotype with the marker indicates that the gene(s) responsible for conferring late blight resistance reside on the chromosome at or near the GO2<sub>586</sub> marker.

Accordingly, the marker can be used as a molecular tag to follow resistance through breeding programs and to identify and isolate disease-resistant breeding stock. Additionally, knowing the relative proximity of the

resistance gene(s) to the marker, isolation of the gene(s) will be facilitated.

The RFLP and RAPD markers CT88 and PO9<sub>587</sub> discussed above can also be used as molecular tags to follow resistance through breeding programs. The use of a combination of the resistance-associated RAPD and RFLP markers can provide an added advantage in following segregation in breeding, and eventually in isolating and cloning the *S. bulbocastanum* gene(s) responsible for conferring resistance to late blight and other diseases.

#### REFERENCES

- Austin, S., M.A. Baer and J.P. Helgeson (1985). Transfer of resistance to potato leaf roll virus from *Solanum brevidens* into *Solanum tuberosum* by somatic fusion. *Plant Science* **39**: 75-82.
- Austin, S., E. Lojkowska, M.K. Ehlenfeldt, A. Kelman and J.P. Helgeson (1988). Fertile interspecific somatic hybrids of *Solanum*: A novel source of resistance to *Erwinia* soft rot. *Phytopathology* **78**: 1216-1220.
- Austin, S., J.D. Pohlman, C.R. Brown, H. Mojtahed, G.S. Santo, D.S. Douches and J.P. Helgeson (1993). Interspecific somatic hybrids between *Solanum tuberosum* and *S. bulbocastanum*: incorporation of nematode resistance. *Am. Potato J.* **70**: 485-495.
- Black, W. and M.E. Gallegly (1957). Screening of *Solanum* species for resistance to physiologic races of *P. infestans*. *Am. Potato J.* **34**: 273-281.
- Bonierbale, M.W., R.L. Plaisted and S.D. Tanksley (1988). RFLP maps based on a common set of clones reveal modes of chromosomal evolution in potato and tomato. *Genetics* **120**: 1095-1103.
- Brown, C.R., C.P. Yang, H. Mojtahedi, G.S. Santo, R. Masuelli (1996). RFLP analysis of resistance to Columbia root-knot nematode derived from *Solanum bulbocastanum* in a BC2 population. *Theor. Appl. Genet.* **92**: 572-576.

- Ehlenfeldt, M.K. and J.P. Helgeson. 1987.  
Fertility of somatic hybrids from protoplast fusions of  
*Solanum brevidens* and *S. tuberosum*. Theor. Appl. Genet.  
73: 395-402.
- 5           Haberlach G., B.A. Cohen, N.A. Reichert, M.A.  
Baer, L.E. Towill and J.P. Helgeson (1985). Isolation,  
culture and regeneration of protoplasts from potato and  
several related *Solanum* species. Plant Sci 39:67-74.
- Helgeson, J.P., G.J. Hunt, G.T. Haberlach and  
10   S. Austin (1986). Somatic hybrids between *Solanum*  
*brevidens* and *Solanum tuberosum*: Expression of a late  
blight resistance gene and potato leaf roll resistance.  
Plant Cell Reports 3: 212-214.
- Helgeson, J.P., G.T. Haberlach, M.K.  
15   Ehlenfeldt, G. Hunt, J.D. Pohlman and S. Austin (1993).  
Fertile somatic hybrids of potato and wild *Solanum*  
species: potential for use in breeding programs. Am.  
Potato. J. 70: 437-452.
- Kim-Lee, H., S.U. Choi, M.S. Chae, S.M. Wielgus  
20   and J.P. Helgeson (1993). Identification of somatic  
hybrids produced by protoplast fusion between *Solanum*  
*commersonii* and *S. tuberosum* haploids. Korean Soc. of  
Plant Tissue Culture 20: 337-344.
- Lam, S.L. (1977). Regeneration of plantlets  
25   from single cells in potatoes. Am.Potato J 54:575-580.
- Lander, E.C., P. Green, J. Abrahamson, A.  
Barlow, M.J. Daly et al. (1987). MAPMAKER: an  
interactive computer package for constructing primary  
genetic linkage maps of experimental and natural  
30   populations. Genomics 1: 174-181.
- McGrath, J.M., S.M. Wielgus, and J.P. Helgeson  
(1994). Recombination and introgression of *Solanum*  
*brevidens* DNA in progeny of a somatic hybrid with potato.  
Amer. Potato J. 71: 686-687
- 35           McGrath, J.M., S.M. Wielgus, and J.P. Helgeson  
(1996). Segregation and recombination of *Solanum*  
*brevidens* syntenic groups in progeny of somatic hybrids

with *S. tuberosum*: Intragenomic equals or exceeds intergenomic recombinations. *Genetics* **142**: 1335-1348.

Novy, R.G. and J.P. Helgeson (1994).

- 5 Resistance to potato virus Y in somatic hybrids between *Solanum etuberosum* and *S. tuberosum* X *S. berthaulii* hybrid. *Theor. Appl. Genet.* **89**: 783-786.

- 10 Tanksley, S.D., M.W. Ganai, J.P. Prince, M.C. DeVicente, M.W. Bonnierbale et al. (1992). High density molecular linkage maps of the tomato and potato genomes. *Genetics* **132**: 1141-1160.

Williams, C.E., G. Hunt, and J.P. Helgeson (1990). Fertile somatic hybrids of *Solanum* species: RFLP analysis of hybrid and its sexual progeny from crosses with potato. *Theor. Appl. Genet.* **80**: 545-551.

- 15 Williams, J.G.K., A.R. Kubelik, K.J. Liwak, J.A. Rafalsi and S.V. Tingley (1990). DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucl. Acids Res.* **18**: 6531-6535.

- 20 The present invention is not limited to the embodiments described and exemplified above, but is capable of variation and modification without departure from the scope of the appended claims.

What is claimed is:

1. A late blight-resistant potato plant comprising a segment of a genome from *Solanum*  
5 *bulbocastanum* which comprises a gene that confers said resistance to late blight.
2. The potato plant of claim 1, wherein the  
10 segment of the *Solanum bulbocastanum* genome is a segment of chromosome 8 of the genome.
3. The potato plant of claim 1, wherein the  
15 gene conferring the late blight resistance co-segregates with a marker selected from the group consisting of a GO2<sub>586</sub> RAPD marker, a PO9<sub>587</sub> RAPD marker, a CT88 RFLP marker, a CT148 RFLP marker, a CT252 RFLP marker and a CT68 RFLP marker.
4. The potato plant of claim 4 wherein the  
20 marker comprises a sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4 AND SEQ ID NO:5.
5. The potato plant of claim 1, which is also  
25 resistant to at least one additional disease selected from the group consisting of potato early blight, *Erwinia* soft rot, and *Verticillium* wilt.
6. The potato plant of claim 1, wherein the  
30 late blight resistance gene is incorporated into the plant by somatic hybridization between a cell of a parent of the plant and a cell of *Solanum bulbocastanum*.
7. The potato plant of claim 1, wherein the  
35 late blight resistance gene is incorporated into the plant by genetic transformation of a cell of the plant with a plant transforming vector comprising the gene.

8. An isolated nucleic acid molecule which is complementary to part or all of a double-stranded molecule selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2 and SEQ ID NO:5.

5

9. The nucleic acid molecule of claim 8, disposed within a vector.

10. A method of monitoring late blight resistance in a breeding cross of progeny of a fertile somatic hybrid of *Solanum tuberosum* and *Solanum bulbocastanum*, which comprises:

- (a) performing the cross;
- (b) isolating genomic DNA from individual progeny of the cross; and
- (c) detecting in the genomic DNA the presence or absence of a genetic marker that is pre-determined to co-segregate with the late blight resistance, the presence or absence of the marker being indicative of the presence or absence of the late blight resistance in the individual progeny of the breeding cross.

11. The method of claim 10, wherein the marker is selected from the group consisting of a GO2<sub>586</sub> RAPD marker, a PO9<sub>587</sub> RAPD marker, a CT88 RFLP marker, a CT148 RFLP marker, a CT252 RFLP marker and a CT68 RFLP marker.

12. A method of identifying a *Solanum bulbocastanum* gene that confers resistance to late blight, said method comprising:

- (a) cloning a DNA segment that co-segregates with the late blight resistance phenotype in progeny of somatic hybrids of *Solanum bulbocastanum* and *Solanum tuberosum*;
- (b) providing a genomic library of the *Solanum bulbocastanum* genome;

(c) isolating clones of the genomic library that contain segments which hybridize with the co-segregating DNA segment; and

(d) identifying at least one gene  
5 disposed within the isolated genomic clones that confer the late blight resistance.


13. The method of claim 12, wherein said  
cloned DNA segment that co-segregates with late blight  
10 resistance comprises part or all of a marker selected from the group consisting of a GO2<sub>586</sub> RAPD marker, a PO9<sub>587</sub> RAPD marker, a CT88 RFLP marker, a CT148 RFLP marker, a CT252 RFLP marker and a CT68 RFLP marker.

15 14. A late blight resistance gene from *Solanum bulbocastanum*, produced by the method of claim 12.

15. A transgenic plant comprising the gene of  
claim 14.


20





*S. bulbocastanum*  
*S. bulbocastanum*  
PI 203900  
PI 203900  
Kathadin  
Kathadin  
Somatic hybrid J101  
Somatic hybrid J138  
Somatic hybrid J138

**FIG. I**

**Figure 2**

3/3




FIG. 3

JUN 26 2000  
 JCS630  
 MARK OFFICE

that my residence, post office address and citizenship are as stated below next to my name;

the specification of which [check one(s) applicable]

X was filed January 24, 2000 as PCT International/U.S. Application No. 09/463,510  
 and was amended by Amendment filed \_\_\_\_\_ (if applicable); [or];  
 is attached to this Declaration, Power of Attorney and Power to Inspect;

that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above; and

that I acknowledge my duty to disclose information which is material to the examination of this application in accordance with Rule 56(a) [37CFR§1.56(a)].

**Prior U.S. Application(s)**

**Filing Date**  
**Day/Mo/Year**

**Status**  
**Pending-Patented-Abandoned**

60/054,267

30 July 1997

Pending

Insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of 35 USC § 112, I acknowledge the duty to disclose material information as defined in Rule 56(a) [37 CFR § 1.56(a)] which occurred between the filing date of the prior U.S. application and the national or PCT international filing date of this application.

**POWER OF ATTORNEY:** As inventor, I hereby appoint **SAUL EWING REMICK & SAUL LLP** of Philadelphia, PA, and the following individual(s) as my attorneys or agents with full power of substitution to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: **Janet E. Reed, Ph.D., Reg. No. 36,252, Alfred W. Zaher, Reg. No. 42,248 and Bruce D. George, Reg. No. 43,631**

**POWER TO INSPECT:** I hereby give **SAUL EWING REMICK & SAUL LLP** of Philadelphia, PA or its duly accredited representatives power to inspect and obtain copies of the papers on file relating to this application.

SEND CORRESPONDENCE TO: SAUL EWING REMICK & SAUL LLP

Centre Square West  
1500 Market Street, 38<sup>th</sup> Floor  
Philadelphia, Pennsylvania 19109

Telephone: (215) 972-7777  
Facsimile: (215) 972-7725

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

**SECOND JOINT INVENTOR (if any)**

Full Name John P. Helgeson  
 First Middle Last  
 Signature [Signature]  
 Date June 8, 2000  
 Residence Madison Wisconsin WI  
 City State or Country  
 Citizenship United States of America  
 Post Office Address:  
809 Oneida Place  
 Madison, Wisconsin 53711  
 City State or Country Zip Code

Full Name Sandra Austin-Phillips  
 Signature Sandra Austin-Phillips  
 Date June 8th, 2000  
 Residence Madison Wisconsin WI  
 City State or Country  
 Citizenship United States of America UK  
 Post Office Address:  
2737 Fox Avenue  
 Madison Wisconsin 53711  
 City State or Country Zip Code

THIRD JOINT INVENTOR (if any)

Full Name Sara K Naess  
First Middle Last

Signature Sara Kristine Naess

Date 6.8.2000

Residence Verona Wisconsin  
City State or Country

Citizenship United States of America NORWAY

Post Office Address:

2586 Prairie Ridge Road

Verona Wisconsin 53593  
City State or Country Zip Code

FOURTH JOINT INVENTOR (if any)

Full Name \_\_\_\_\_  
First Middle Last

Signature \_\_\_\_\_

Date \_\_\_\_\_

Residence \_\_\_\_\_  
City State or Country

Citizenship \_\_\_\_\_

Post Office Address:

\_\_\_\_\_  
City State or Country Zip Code

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- 5 (i) APPLICANT: Helgeson, John P.  
Austin, Sandra  
Naess, Sara K.
- 10 (ii) TITLE OF THE INVENTION: GERMPLASM AND MOLECULAR  
MARKERS FOR DISEASE RESISTANCE IN POTATO
- (iii) NUMBER OF SEQUENCES: 5
- 15 (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: Dann, Dorfman, Herrell & Skillman  
(B) STREET: 1601 Market Street, Suite 720  
(C) CITY: Philadelphia  
(D) STATE: PA  
(E) COUNTRY: USA  
20 (F) ZIP: 19103
- (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Diskette  
(B) COMPUTER: IBM Compatible  
25 (C) OPERATING SYSTEM: DOS  
(D) SOFTWARE: FastSEQ Version 1.5
- (vi) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER: Not yet assigned  
30 (B) FILING DATE: 27-JUL-1998
- (vii) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER: US 60/054,267  
35 (B) FILING DATE: 30-JUL-1997
- (viii) ATTORNEY/AGENT INFORMATION:  
(A) NAME: Janet E. Reed  
(B) REGISTRATION NUMBER: 36,252  
(C) REFERENCE/DOCKET NUMBER: P98003WO  
40
- (ix) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: 215-563-4100  
(B) TELEFAX: 215-563-4044
- 45 (2) INFORMATION FOR SEQ ID NO:1:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 586 base pairs  
(B) TYPE: nucleic acid  
50 (C) STRANDEDNESS: double  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA  
(iii) HYPOTHETICAL: NO  
55 (iv) ANTISENSE: NO  
(v) FRAGMENT TYPE:  
(vi) ORIGINAL SOURCE: Solanum bulbocastanum

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	GGCACTGAGG	GGTAGTAAGC	CTCCTGCATG	TACTAAGTAT	GGTAGATCCA	CTCAGGGTTG	60
	TGCCATGATG	GCTTAAGTGG	TTGTTTCAAG	TATGGCCAGA	ACGGTTATTT	TATGAGAGAG	120
5	TTTCCAAAGA	ACATGCAGGG	TAATGGTAAT	GGGGATAATA	GAACCCAGTC	TTCTTCAGTG	180
	ACTCCACCAG	ACAGAGCTGC	ATCTAGAGGA	GCTACTTCGA	GGCAGGCGGA	GGATCGAACC	240
	TCTTTATGCT	ATCACTAGTC	GCCAAGAGAA	AGAGGATTCG	CCAGATGTTG	TCACTGGTAT	300
	GATCCAAGTC	TTTAACTTTG	ATTTTATACT	TTTCTAGATC	CAGGAGCGAG	TTTATCCTTT	360
	GTAACCTCTT	ATGTTGCGGT	TAATTTTGAT	GTTCTTCCTA	AGAACTTAT	TGAGCCCTTC	420
10	AGTGTCTTCTA	CACTTGTGTTG	TCTATTATAG	TAGAGAGAGT	CTGTTATGAT	TGTACCGTTT	480
	TCGTCAATCA	CAAGAGCACC	ATGGTTGATT	TAGTTGAGTT	AGACATGGTA	GAATTTGATG	540
	TTATTCTTGG	TATGGACTGA	CTTCATTCTT	GTTATGCCTC	AGTGCC		586

## (2) INFORMATION FOR SEQ ID NO:2:

15

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 587 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

20

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (iv) ANTISENSE: NO

25

## (v) FRAGMENT TYPE:

## (vi) ORIGINAL SOURCE: Solanum bulbocastanum

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

30	GTGGTCCGCA	TATAACTCAA	GAAGTTGTAA	TGCATGTATC	GGATATATGT	ATACATGTTG	60
	TCTTTTGCAA	AGTTTACTTT	TTTATTANTT	AATCTTGTTT	GTGTCTGGAG	GTGGTGGTGG	120
	GGTGGGATAG	TGGTGAAGCT	AGAAATTTAG	TTAAGTGTGT	TCAAGATTTA	AATATACATA	180
	TGAAAAATAA	TTTTTGATCT	ATATATATAG	TTATAATTTT	NTGATGAAGG	TAGTTCAACT	240
	GACCACCCGT	NACTACATGT	GGCTACCGTA	CTGGGTGGGG	TGGAAGGTCN	TGGTGTTTAC	300
35	AGTGATGTGG	GGGCCTCTGA	AATGCTTTTG	TGGGCAATGT	GGGAATTACT	GTTTATCTTT	360
	TCTTTATTGA	AGTCATTGAG	TGTTTGAGTT	ATTTAACTAT	GAAAGGTAGC	TAGTGGGTAA	420
	TGTTATTGAT	GACTTTGTGT	GAAGAACAAA	ATGTCAATCA	TTCAGAGCGT	TCAATGGGGG	480
	CACGTGCTGG	AGTCCCATT	GGATTATTGG	GTTTAGGGGT	TGCCTAGGTG	GTGGTGGTGG	540
	TGGTTGGTAA	TGGATAATAA	TGTTGGTGAA	ATATGGGTGC	GGAGGAG		587

40

## (2) INFORMATION FOR SEQ ID NO:3:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 596 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: linear

45

## (ii) MOLECULE TYPE: cDNA

## (iii) HYPOTHETICAL: NO

## (iv) ANTISENSE: NO

## (v) FRAGMENT TYPE:

## (vi) ORIGINAL SOURCE: tomato

55

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

	GTTGGGCAGA	AGAGCTAGGA	AGAGTAAGCA	TGTCAAGTGA	TAGTTGCAGC	CACTGGTGT	60
	ATAGTTGTAG	ACAACCCGTG	AATCTCAGGA	GACAAAATGA	TGTTTGCCCC	AATTGCGGTG	120
	GTGGATTTGT	TCAAGAGCTT	GAAGACATAA	CGAGTAGTAG	TCTAGATAAT	CAGACCCAGA	180
60	GGCCGAGATT	CATGGAATCC	GTCTCAAACT	TTTAAAGACG	ACAAATCTCA	GCTACAAGTA	240
	ATACTTCTGA	GAGAGGGAGA	TCTGATGGGG	GTGCTGAACG	AGGAAATTTG	TGGAATCCGT	300

	TGCTGATTTT	CAGTGGTGAT	ACGCCTGTTC	ATATGCCTGG	GGATGGTGGA	GTTTTGGAGT	360
	TTCTTAATGA	GGCACCTTGG	ATTCCGAGCA	AGAAAATGGT	GGTGATTATT	TTGTTGGTCC	420
	AGGAGTGGAG	GAATTTTTTG	AAGAAATGT	AAATAGAAAT	CAGCGTGGTG	CTCCTCCTCC	480
	TGTCTCGAGA	TGTTCAATTG	ATTCCCTACC	AACAGTCAAG	ATATCAAAAA	AGGATGTTAG	540
5	ATCGGATTCT	CACTGCCCTG	TTTGTAAGGA	GAAATTTGCT	CTGGGGACTA	AGGCAA	596

## (2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
- 10 (A) LENGTH: 589 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
(D) TOPOLOGY: linear
- 15 (ii) MOLECULE TYPE: cDNA  
(iii) HYPOTHETICAL: NO  
(iv) ANTISENSE: NO  
(v) FRAGMENT TYPE:  
(vi) ORIGINAL SOURCE: Solanum tuberosum
- 20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

	GTTGGGCAGA	AGAGCTAGGA	AGAGTAAGCA	TGTCAAGTGA	CAGTTGCAGC	CACTGGTGTT	60
	ATAGTTGTAG	ACAACCCGTG	AATCTCAGGA	GACAAAATGA	TGTTTGCCCC	AATTGCAGTG	120
25	GTGGATTGT	TCAAGAGCTT	GAAGACATAA	CGAGTAGTAG	TGTAGATAAT	CAGAGCCAGA	180
	GGCCGAGATT	TATGGAATCC	GTCTCAAAC	TTTAAAGACG	ACAAATCGCT	ACAAGTAATA	240
	CTTCTGAGAG	AGGGAGATCT	GATGGGGGTG	CTGAACGAGG	AAATTTATGG	AATCCATTGC	300
	TGATTTTCAG	TGGTGATACG	CCTGTTCAAG	ATGCCTGGGG	ATGGTGAGT	TTTGGAGTTT	360
	CTTAATGAGG	CTCTTGGCTT	CCGACAAGAA	AATGGTGGTG	ATTATTTTGT	TGGTCCAGGA	420
30	GTGGAGGAAT	TTTTTGAAGA	AATTGTAAAT	AGAAATCAGC	GTGGTGCTCC	TCCTGCCTCA	480
	AGATGTTCAA	TTGATTCCT	ACCAACAGTC	AAGATATCGA	AAAAAGATGT	TAGATCGGAT	540
	TCTCACTGCC	CTGTTTGTA	AGAGAAATTT	GCTCTGGGGA	CTAAGGCAA		589

## (2) INFORMATION FOR SEQ ID NO:5:

- 35 (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 592 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: double  
40 (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA  
(iii) HYPOTHETICAL: NO  
(iv) ANTISENSE: NO  
45 (v) FRAGMENT TYPE:  
(vi) ORIGINAL SOURCE: Solanum bulbocastanum

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

50	GTTGGGCAGA	AGAGCTAGGA	AGAGTAAGCA	TGTCAAGTGA	CAGTTGCAGC	CACTGGTGTT	60
	ATAGTTGTAG	ACAACCCGTG	AATCTCAGCA	GACAAAATGA	TGTTTGCCCC	AATTGCGGTG	120
	GTGGATTGT	TCAAGAGCTT	GAAGACATAA	CGAGTAGTAG	TGTAGATAAT	CAGAGCCAGA	180
	GGCCGAGATT	CATGGAATCC	GTCTCAAAC	TTTAAAGACG	ACAAATCCCA	ACTACAAGTA	240
	ATACTTCTTG	AGAGAGGGAG	ATCTGATGGG	GGTGCTGAAC	GAGGAAATTT	GTGGAATCCG	300
55	TTGCTGATT	TCAGTGGTGA	TACACCTGTT	CGGATCGCTG	GGGATGGTGG	AGTTTGGAG	360
	TTTCTTAATG	AGGCTCTTGG	CTTTCGACAA	GAAAAAGGTG	GTGANTATTT	TGTTGGCCCA	420
	GGAGTGGAGG	AGTTTTTTGA	AGAAATTGTA	AATAAAAATC	AGCGTGGTGC	TCCTCCTGTC	480
	TCAAGATGCT	CAATTGATTC	CCTACCAACA	GTCAAGATAT	CGAAAAAGGA	TGTTAGATCG	540
	GATTCTCACT	GCCCTGTTTG	TAAAGAGAAA	TTTGCTCTGG	GGACTAAGGC	AA	592